NOTE ON SOME HOLOCENE MARINE SHELLS FROM THE ARAN ISLES, CO. GALWAY.

By J. R. LE B. TOMLIN, M.A., F.E.S.

Read 14th April, 1916.

Mr. Kennard sent me, some little time ago, a sample of blown sand of Holocene age which he had received from Mr. R. A. Phillips, of Cork. The latter writes that he collected it from a low-lying, sandy marsh at Killeany Bay, Inishmore, Aran Isles, Galway, where there was blown sand covered with grass, and the material was actually obtained from a pit excavated by the wind in this ground.

For convenience of reference the names are given according to the last Conchological Society's list (Journ. Conch., vol. x, 1901, pp. 10-26). The remains of bivalves are in such fragmentary condition that it is only possible to identify them in one or two

eases. The following is the list of species:

Mytilus edulis, L., several.
Cardium sp. (?), fragments common
Saxicara arctica, L., one valve.
Acmæa virginea (Müll.), two.
Gibbula cineraria (L.), two.
Phasianella pullus (L.), common.
Lacuna divaricata (F.), one.
L. parva (da C.), one.
Littorina rudis (Maton), several.
Rissoa parva (da C.) and var.
interrupta, Ad., both equally common.

Alvania punctura (Mont.), two. Manzonia costata (J. Ad.), three. Onoba striata (J. Ad.), very common.

— var. aculeus, Gld., one. Cingula semistriata (Mont.), three. Barleeia rubra (Mont.), one. Bittium reticulatum (da C.), com-

Triforis perversa (L.), one. Cerithiopsis concatenata, Conti, one.

Odostomia turrita, Hanley, one. Brachystomia rissoides (Hanley), one.

Pyrgulina interstincta (Mont.), three.

Eulima bilineata (Alder), one. Cæcum imperforatum (G. Ad.), four.

C. glabrum (Mont.), one.
Buccinum undatum, L., one.
Purpura lapillus, I., one.
Mangilia costata (Don.), one.
Clathurella linearis (Mont.), two.
Tornatina truncatula (Brug.),
three.